Worksheet

Embryonic Development

Background Information


The development of the embryo begins after fertilization. It follows a well-defined pattern of growth and **differentiation**. Embryonic development is controlled, for the most part, by the genes - the hereditary material contributed by both parents. Development may occur either in the external environment or within the body of a parent.

Stages of Embryonic Development

After fertilization, the **zygote** divides by mitosis into two cells. Mitotic division, which is called **cleavage** in embryos, continues, eventually forming a solid ball of cells. This is the **morula** stage. As cleavage continues, the center of the ball becomes hollow. This is the **blastula** stage. With still further cleavage, an inward growth of cells at a particular point results in the formation of a pocket, which pushes into the hollow center of the ball. This is the **gastrula** stage.

Within the double-walled gastrula three distinct layers of embryonic tissue - the **germ layers** - develop. The inner layer is the **endoderm**. The outer wall of the gastrula forms the **ectoderm**. And between the ectoderm and endoderm a middle layer called the **mesoderm** forms.

1. Label each stage and structure of embryonic development shown in the following diagrams.



·	, and
Early cell division in embryonic development is commonly called	and can be either
cleavage or	cleavage depending on the animal.
What does the blastopore become in most animals?	What does it become in humans,
other chordates, and echinoderms?	
The embryonic germ layers give rise to all parts of the animal's body. embryonic germ layers:	List a few of the systems that come from the followi
a) ectoderm -	
b) mesoderm -	
c) endoderm -	
Define differentiation.	
Use the following terms to complete the sentence below: specializati	on, multicellularity, and differentiation.